



**LAND MANAGEMENT, ENVIRONMENT & SOLID-WASTE:
INSIDE EDUCATION AND BUSINESS IN CENTRAL ASIA
ERASMUS-EDU-2023-CBHE Project Number: 101129032**

MICRO-CREDENTIAL

**CIRCULAR ECONOMY FOR LAND AND WASTE
MANAGEMENT**

9.- Circular Economy in Agriculture and Rural Areas

Circular Agriculture and Rural Land Systems

PDF W09



Co-funded by
the European Union

Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Education and Culture Executive Agency (EACEA). Neither the European Union nor EACEA can be held responsible for them.

INDEX

1. SUMMARY OF KEY IDEAS	4
2. LEARNING OBJECTIVES.....	4
3. INTRODUCTION.....	5
4. FROM LINEAR AGRICULTURE TO CIRCULAR LAND SYSTEMS	7
4.1. The linear agricultural model	7
4.2. What makes agriculture circular?.....	7
4.3. Systems thinking and territorial circularity	9
5. CLOSING NUTRIENT AND ORGANIC-MATTER LOOPS	9
5.1. Nitrogen, phosphorus and carbon.....	9
5.2. Crop residues, manure and organic waste	11
5.3. Composting and anaerobic digestion	12
6. SOIL HEALTH AND REGENERATIVE LAND MANAGEMENT	14
6.1. Soil degradation as a circularity failure	14
6.2. Regenerative agricultural practices	14
6.3. Circular agriculture in saline and degraded lands	15
7. WATER, ENERGY AND MULTI-USE RURAL SYSTEMS	16
7.1. Circular water management.....	16
7.2. Agricultural biomass and renewable energy	17
7.3. Agrivoltaics and multiple land uses	18
8. CIRCULAR VALUE CHAINS AND RURAL BIOECONOMY	18
8.1. From agricultural residues to secondary resources	18
8.2. Biomass cascading	19
8.3. Industrial symbiosis in rural territories	19
9. CIRCULAR AGRICULTURE IN CENTRAL ASIA	20
9.1. Regional challenges	20
9.2. Circular opportunities.....	21
9.3. A practical transition framework.....	23
10. GOVERNANCE, ECONOMICS AND ENABLING CONDITIONS	24

11.	<i>MONITORING CIRCULAR PERFORMANCE</i>	25
12.	<i>CONCLUSIONS</i>	27
13.	<i>REFERENCES</i>	28

1. SUMMARY OF KEY IDEAS

Circular agriculture applies circular economy principles to food production, agricultural land, biomass, nutrients, water and rural territories. Instead of following a linear model in which external inputs are consumed, products leave the farm and residues become waste, circular agriculture seeks to maintain resources in productive use for as long as possible and return biological nutrients safely to soil.

The concept goes beyond recycling agricultural waste. A truly circular agricultural system seeks to reduce unnecessary inputs, prevent nutrient losses, restore soil functions, conserve water, recover organic matter and energy, integrate crop and livestock activities, and create connections among farms, food processors, municipalities, waste managers and other rural actors.

At field scale, circularity can involve crop rotations, legumes, cover crops, reduced tillage, compost, digestate, precision fertilisation and efficient irrigation. At farm scale, it can integrate livestock manure, crop residues, energy production and nutrient management. At territorial scale, it may connect farms with food industries, cities, wastewater treatment plants, biogas facilities and markets for secondary materials.

This territorial dimension is particularly important. A farm cannot necessarily close every loop internally. A crop farm may need organic nutrients from a livestock facility; a food-processing plant may send by-products to animal-feed production or anaerobic digestion; a municipality may provide source-separated organic material for controlled composting; and renewable energy may be shared among rural enterprises.

For Central Asia, circular agriculture has particular relevance because agricultural production takes place under conditions of water scarcity, soil salinisation, land degradation, ageing irrigation infrastructure and significant losses of biomass and nutrients. The transition towards circular rural systems therefore needs to focus on water productivity, soil organic matter, salinity management, biomass recovery, safe nutrient recycling and locally appropriate technologies.

The core message of this lesson is that circular agriculture is not simply about transforming waste into a resource. It is about redesigning agricultural and rural systems so that waste generation, resource loss and land degradation are progressively prevented by design.

2. LEARNING OBJECTIVES

By the end of this lesson, the student will be able to:

- Explain the difference between linear and circular agricultural systems.
- Describe the main principles of circular agriculture.
- Analyse nutrient, organic-matter, water and biomass flows within rural systems.

- Explain the importance of soil health for circular agriculture.
- Compare composting and anaerobic digestion as organic-resource recovery options.
- Apply the biomass cascading principle to agricultural residues.
- Identify opportunities for industrial symbiosis in rural and agri-food systems.
- Evaluate circular agriculture options for arid and semi-arid regions.
- Identify specific challenges and opportunities for circular agriculture in Central Asia.
- Propose indicators for monitoring circularity at farm and territorial scales.

3. INTRODUCTION

Agriculture is fundamentally dependent on cycles. Plants take nutrients from soil, animals consume biomass, organic residues decompose, water circulates through soil and vegetation, and carbon moves among the atmosphere, plants and soil. Natural ecosystems operate largely through such cycles, with material continuously transformed and reused.

Modern agricultural systems have often weakened these connections. Nutrients are imported as mineral fertilisers, crops are harvested and transported over long distances, organic residues may be burned or discarded, manure may accumulate in areas with insufficient land for safe application, and food waste is separated from the farms that originally produced the food.

This creates what can be understood as a **linear agricultural metabolism**. Resources enter the system, products leave it, and a significant proportion of nutrients, biomass, water and energy is lost along the way.

The environmental consequences include soil degradation, nitrate leaching, eutrophication, greenhouse-gas emissions, groundwater contamination, loss of organic matter, inefficient water use and increasing dependence on external resources.

Circular agriculture seeks to change this logic.

It asks a different set of questions:

Where do nutrients come from? Where do they go after harvest? Can organic residues remain productively within the system? Can manure replace part of mineral fertiliser demand? Can wastewater be safely treated and reused? Can food-processing by-products become animal feed, biochemical products or energy? Can one farm's surplus become another farm's resource?

As illustrated conceptually in Figure 1, the transition from linear to circular agriculture replaces one-way flows with managed loops involving nutrients, biomass, water, energy and information.




Figure 1. From linear agriculture to circular agriculture

Did you know..?

Circular agriculture does not require every individual farm to be completely self-sufficient. Some resource loops can only be closed effectively at landscape or regional scale through cooperation among crop farms, livestock farms, food processors, municipalities, water managers and waste-treatment facilities.

4. FROM LINEAR AGRICULTURE TO CIRCULAR LAND SYSTEMS

4.1. The linear agricultural model

The conventional linear model can be summarised as **take–produce–consume–discard**.

At farm level, inputs may include mineral fertilisers, pesticides, irrigation water, animal feed, fuel, plastics and machinery. Crops and animal products are then exported from the farm. With them leave nitrogen, phosphorus, carbon and other nutrients.

If nutrients and organic matter are not returned in a controlled way, soil fertility becomes increasingly dependent on external inputs.

The problem is not agricultural production itself. Modern agriculture has achieved enormous increases in productivity. The problem is a system in which resource use and production are disconnected from the management of residual materials and environmental impacts.

Consider a simplified cereal system. Mineral nitrogen is applied to the soil. Part is absorbed by the crop, but another part may be lost by nitrate leaching, ammonia volatilisation, surface runoff or gaseous emissions. The grain is harvested and transported away. Straw may be removed or burned. Little organic matter returns to the soil.

The result is not a closed nutrient cycle but a sequence of inputs and losses.

The same linear logic can occur with water. Water is diverted from a river or pumped from an aquifer, transported through inefficient canals, applied to fields and lost through evaporation, uncontrolled drainage or deep percolation. In saline environments, poor water and drainage management can also transport salts and create long-term soil degradation.

Agricultural plastics provide another example. Mulch films, pesticide containers, irrigation pipes and packaging may improve productivity but become a pollution source if no collection and recovery system exists.

Linear agriculture therefore produces more than food. It can also generate nutrient surpluses, polluted water, greenhouse gases, degraded soil, waste biomass and abandoned materials.

4.2. What makes agriculture circular?

Circular agriculture is based on the deliberate management of resources as cycles rather than one-way flows.

Five principles are particularly important.

First, nutrient loop closure. Nitrogen, phosphorus and other nutrients should be used efficiently and recovered where technically safe and economically feasible.

Second, soil regeneration. Soil should be treated as living natural capital rather than merely as a physical substrate. Maintaining organic matter, biological activity, structure and water-holding capacity is fundamental.

Third, cascading use of biomass. Biomass should normally be directed to the highest-value feasible use before lower-value recovery. Food comes before feed; feed may come before materials; materials may come before energy; and nutrients should ultimately be returned safely to land.

Fourth, diversity and integration. Crop rotations, livestock integration, agroforestry, biodiversity and complementary enterprises can create internal synergies and reduce dependence on external inputs.

Fifth, territorial collaboration. Circularity often requires connections among different actors. A single farm may produce too little organic waste for its own biogas plant, but several farms together may support a collective facility.

These principles are summarised in Table 1.

Table 1. Core principles of circular agriculture

Principle	Main idea	Practical agricultural application
Nutrient loop closure	Reduce nutrient losses and recover useful N, P and organic matter	Compost, digestate, manure management, nutrient budgeting
Soil regeneration	Maintain and restore the biological and physical functions of soil	Cover crops, rotations, reduced tillage, organic amendments
Biomass cascading	Use biomass first for the highest-value suitable purpose	Food → feed → materials → energy → nutrient return
Integration and diversity	Link complementary biological and productive processes	Crop–livestock integration, agroforestry, legumes
Territorial circularity	Connect resource producers and users across a rural region	Industrial symbiosis, collective biogas, regional composting

As shown in Table 1, circular agriculture combines biological processes with technological, organisational and territorial changes. It should therefore not be confused with any single farming method.

Organic farming, regenerative agriculture, agroecology, precision agriculture and conservation agriculture may all contribute to circularity, but none is automatically equivalent to circular agriculture. The key question is how successfully the system reduces virgin inputs, losses and waste while maintaining productivity and regenerating natural capital.

**Reflect**

A farm replaces mineral fertiliser with compost transported 800 km by truck. Is this necessarily a more circular solution? Consider nutrient recovery, transport impacts, product quality and the possibility of finding a closer source.

4.3. Systems thinking and territorial circularity

A circular farm should not be analysed in isolation.

Suppose a dairy farm produces manure that exceeds the nutrient absorption capacity of its own land. Applying all manure locally could create nitrate and phosphorus pollution. From a farm-level perspective, the manure is a surplus. At territorial level, however, nearby crop farms may have a nutrient deficit.

The circular opportunity is not to apply more manure to the dairy farm's own fields. It is to connect nutrient surplus and nutrient demand.

The same logic applies to agricultural residues. A fruit-processing plant may produce pomace that becomes a waste-disposal problem. Depending on quality and composition, it could instead be processed into pectin, used as livestock feed, digested to produce biogas or composted.

Circularity is therefore spatial. Transport distance, timing, quality, infrastructure and market demand determine whether loops can be closed in practice.

This leads to a key principle:

The smallest possible loop is not always the best loop. The appropriate scale is the one that achieves safe, efficient and environmentally beneficial resource circulation.

5. CLOSING NUTRIENT AND ORGANIC-MATTER LOOPS

5.1. Nitrogen, phosphorus and carbon

Nutrient management lies at the centre of circular agriculture.

Nitrogen is essential for crop productivity but highly reactive and mobile. When nitrogen inputs exceed crop uptake or are poorly timed, losses can occur through nitrate leaching, ammonia volatilisation, runoff and gaseous emissions.

Circular nitrogen management seeks first to improve efficiency. The objective is not merely to recycle more nitrogen but to avoid unnecessary surplus.

This can involve:

- matching fertiliser application to crop demand;
- using legumes and crop rotations;
- recovering nitrogen from manure and organic residues;
- improving manure storage;
- using precision application technologies;
- preventing nitrate leaching during fallow periods.

Phosphorus presents a different challenge. Unlike nitrogen, it has no large atmospheric reservoir that can be biologically fixed. Agricultural phosphorus is largely derived from phosphate rock, and phosphorus losses can contribute to eutrophication.

Circular phosphorus management therefore places particular emphasis on recovery from manure, wastewater, sewage sludge and other organic streams, provided contaminant risks are controlled.

Carbon connects circular agriculture directly to soil health. Crop residues, roots, manure, compost and other organic materials can contribute to soil organic matter. Soil organic carbon supports aggregation, biological activity, nutrient cycling, erosion resistance and water retention.

However, simply applying organic residues does not guarantee soil carbon accumulation. The result depends on climate, soil type, decomposition rate, residue quality, tillage and long-term management.

The interactions among nitrogen, phosphorus and carbon are represented in Figure 2.




Figure 2. Circular nutrient and biomass flows in an integrated rural system

5.2. Crop residues, manure and organic waste

Agricultural residues are often described as waste, but their role is more complex.

Crop residues may provide several functions:

- maintaining soil organic matter
- protecting soil against erosion
- conserving moisture
- feeding livestock
- providing raw material for bio-based products
- producing energy

These uses can compete with each other.

For example, removing all cereal straw for biogas or combustion may generate renewable energy but reduce soil organic-matter inputs. Leaving all residues in the field may support soil health but miss opportunities for higher-value uses.

Circular management therefore requires balance.

The biomass cascade provides a useful framework. Edible biomass should normally remain food. Suitable food-processing by-products may be used as feed. Fibres and compounds may be extracted for materials. Energy recovery generally comes later in the cascade.

Manure also illustrates the need for careful management. It contains valuable nitrogen, phosphorus, potassium and organic matter, but uncontrolled application can pollute water and generate emissions.

A circular manure strategy should answer four questions:

- What nutrients does the manure contain?
- What nutrients do crops require?
- When and where should it be applied?
- Can treatment improve storage, transport or nutrient recovery?

The goal is not maximum recycling. It is safe and efficient nutrient substitution.



An organic residue does not become environmentally safe merely because it is called a circular resource. Digestate, compost, manure, sewage-derived products and other recycled amendments must be evaluated for nutrients, salinity, pathogens, heavy metals, persistent contaminants and physical impurities where relevant.

5.3. Composting and anaerobic digestion

Composting and anaerobic digestion are two important technologies for closing organic-resource loops.

Composting is an aerobic biological process. Microorganisms transform organic materials under controlled moisture, aeration and temperature conditions into a more stable amendment.

Good compost can return organic matter and nutrients to soil and improve structure. However, poor source separation can introduce plastics, glass, metals or other contaminants.

Anaerobic digestion, by contrast, occurs without oxygen. Organic material is transformed into biogas and digestate. The biogas can be used for heat, electricity, upgraded biomethane or other energy purposes.

Digestate retains much of the original nutrient content and may be used as fertiliser when quality and agronomic conditions are suitable.

The comparison is shown in Table 2.

Table 2. Comparison of composting and anaerobic digestion

Criterion	Composting	Anaerobic digestion
Biological environment	Aerobic	Anaerobic
Main output	Stable organic amendment	Biogas and digestate
Energy recovery	No direct net energy production	Renewable gas production
Organic matter	Produces stabilised compost	Digestate retains partially stabilised organic matter
Nutrient management	Some nitrogen may be lost during processing	Much nitrogen remains in digestate
Main advantages	Simpler technology, soil organic-matter benefits	Energy plus nutrient recovery
Main limitations	Odour, moisture, contamination and N losses if poorly managed	Higher capital needs and careful digestate management
Typical suitable feedstocks	Green waste, crop residues, source-separated organics	Manure, food waste, wet organic residues

As shown in Table 2, neither technology is universally superior. Composting may be more suitable for dry lignocellulosic residues, while anaerobic digestion can be particularly attractive for wet, easily biodegradable organic streams.

In some systems, both can be combined. Organic material can first undergo anaerobic digestion to recover energy, after which the solid fraction of digestate may be composted to produce a more stable soil amendment.



A rural district produces cattle manure, fruit-processing residues, wheat straw and municipal food waste. Decide which materials could be suitable for composting, anaerobic digestion, direct soil return or higher-value use. Explain why one single treatment option would probably not be optimal for all four streams.

6. SOIL HEALTH AND REGENERATIVE LAND MANAGEMENT

6.1. Soil degradation as a circularity failure

Healthy soil is the foundation of circular agriculture.

A system cannot be called circular if it recovers materials while degrading the land on which production depends. Soil erosion, organic-matter loss, compaction, salinisation, contamination and biodiversity decline all represent losses of natural capital.

A degraded soil becomes more dependent on external inputs. Poor structure can reduce infiltration. Low organic matter can decrease water retention. Erosion removes nutrients. Salinisation limits plant uptake. Compaction restricts roots.

These changes create a negative feedback loop:

degradation → lower soil function → greater external input dependence → further environmental pressure.

Circular soil management seeks the opposite:

organic matter recovery → better soil structure → improved nutrient and water efficiency → greater resilience → reduced losses.

This does not mean that organic amendments alone can repair every degraded soil. Salinity, sodicity, erosion, contamination and compaction require specific diagnoses and interventions.

6.2. Regenerative agricultural practices

Several practices can support both soil health and circularity.

Crop rotations diversify nutrient demand and biological processes. Including legumes can reduce dependence on synthetic nitrogen.

Cover crops protect soil between commercial crops, reduce erosion and may capture residual nitrogen that would otherwise be lost.

Reduced tillage, where agronomically appropriate, can reduce physical disturbance and erosion. Its effects on carbon storage depend strongly on local conditions and soil depth.

Organic amendments can return carbon and nutrients. Their use should be based on quality control and agronomic demand.

Agroforestry integrates woody vegetation with crops or livestock and can provide biomass, shelter, microclimate regulation, erosion protection and biodiversity benefits.

Precision agriculture can also support circularity by reducing over-application and matching nutrients or irrigation more closely to spatially variable crop needs.

Circular agriculture is therefore not necessarily low-tech or high-tech. It may combine traditional biological knowledge with sensors, remote sensing, automated irrigation and digital nutrient accounting.



Many practices now described as circular or regenerative—crop rotations, nutrient return, mixed crop–livestock systems and reuse of organic residues—have historical precedents extending back centuries. The innovation lies partly in combining this knowledge with modern monitoring, quality control and territorial planning.

6.3. Circular agriculture in saline and degraded lands

Circular agriculture in arid and semi-arid regions must address different priorities from those in humid regions.

Water scarcity changes everything.

Organic residues may improve soil structure and water retention, but their salinity must be controlled. Reusing drainage water may conserve freshwater but increase salt loading. Applying manure or digestate can return nutrients but may worsen salinity or nitrate pollution if overused.

Central Asian agricultural landscapes frequently require integration of:

- irrigation efficiency
- drainage
- salinity control
- soil organic-matter restoration
- crop diversification
- nutrient budgeting
- safe biomass recovery

The key is not to maximise each individual loop separately. A system that maximises water reuse but progressively accumulates salts is not circular in a sustainable sense.



Reflect

Consider an irrigated farm where drainage water is collected and reused. Water withdrawals decrease, but soil salinity increases every year. Is this a successful circular system? What additional indicators would you need before deciding?

7. WATER, ENERGY AND MULTI-USE RURAL SYSTEMS

7.1. Circular water management

Agriculture is the dominant water user in many arid and semi-arid regions. Circular water management involves two complementary strategies:

use less water per unit of useful production, and recover or reuse water safely where appropriate.

Efficiency measures can include improved conveyance, drip irrigation, micro-irrigation, soil-moisture monitoring, irrigation scheduling and crop selection.

Reuse options can include treated municipal wastewater, reclaimed agro-industrial water, drainage water and captured rainwater.

However, water reuse requires careful attention to water quality.

Relevant concerns may include:

- salinity
- sodium
- nutrients
- pathogens
- trace elements
- organic pollutants
- pesticides

The suitability of reclaimed water depends on the crop, soil, irrigation method, exposure pathways and local standards.

For Central Asia, circular water management must be understood at basin scale as well as farm scale. The Aral Sea crisis demonstrated the consequences of large-scale water diversion, inefficient irrigation and insufficient consideration of environmental flows.

Saving water at field level does not automatically guarantee basin-level water savings. For example, water previously considered a "loss" through seepage may recharge groundwater or support downstream users.

Circular water management therefore requires hydrological systems thinking.



Higher irrigation efficiency at farm level does not necessarily produce an equivalent reduction in total basin water consumption. Return flows, groundwater recharge and changes in cropped area must also be considered.

7.2. Agricultural biomass and renewable energy

Agricultural systems produce significant biomass streams, including straw, stalks, husks, pruning residues, manure and food-processing residues.

Some can support renewable energy production through:

- anaerobic digestion
- combustion
- gasification
- fermentation
- production of biofuels

But circular bioenergy should follow the cascade principle.

Burning a high-quality biomass resource that could first serve as food, feed or material may destroy value. Conversely, anaerobic digestion of unavoidable manure or food-processing waste can recover energy while preserving nutrients in digestate.

Residue removal also has ecological limits. Crop residues protect soil and contribute to organic-matter cycling. Excessive extraction may increase erosion or reduce soil carbon.

Therefore, the circular question is not:

How much biomass can be removed?

It is:

How much can be used for additional applications after maintaining essential soil and ecosystem functions?

7.3. Agrivoltaics and multiple land uses

Agrivoltaics combines agricultural production and photovoltaic electricity generation on the same land.

Solar panels may be elevated above crops or arranged to permit grazing or cultivation between rows. Under suitable designs, partial shading may reduce heat stress and evapotranspiration for some crops.

Agrivoltaics illustrates a broader circular principle: **multiple functions from the same land resource**.

However, good design is essential. Poorly planned systems can restrict machinery, reduce crop yields or become primarily energy installations with agriculture as a secondary justification.

The objective should be to optimise combined land productivity rather than simply place two uses together.

8. CIRCULAR VALUE CHAINS AND RURAL BIOECONOMY

8.1. From agricultural residues to secondary resources

Circular agriculture extends beyond the farm gate.

Agricultural products move through processing, distribution and consumption systems. Along the way, they generate by-products such as:

- cereal bran
- fruit pomace
- whey
- spent grain
- oilseed residues
- slaughterhouse by-products
- food waste

In a linear system, these materials become disposal problems.

In a circular value chain, they are characterised, separated and directed towards useful applications.

A brewery can provide spent grain for animal feed. A fruit processor can recover pectin or send residues to digestion or composting. A dairy plant can valorise whey. Food residues can produce biogas.

The feasibility depends on composition, quality, regulation, quantity, distance and market demand.

8.2. Biomass cascading

The cascading principle seeks to retain the highest possible value of biomass.

A simplified hierarchy is:

human food → animal feed → bio-based products → biochemical extraction → energy → nutrient return to soil

This sequence is not absolute, but it prevents a common circularity mistake: using resources immediately for low-value energy recovery when higher-value options remain feasible.

For example, edible food should normally not be converted to biogas merely because digestion is technically possible. Prevention and redistribution preserve more value.

Likewise, a processing by-product suitable for feed may provide greater resource efficiency than direct combustion.

8.3. Industrial symbiosis in rural territories

Industrial symbiosis occurs when the output or by-product of one activity becomes a useful input to another.

In rural systems, examples include:

- brewery residues → livestock feed
- fruit-processing residues → pectin, compost or biogas
- manure → anaerobic digestion → biogas and digestate
- food-industry waste heat → greenhouses
- municipal organic waste → controlled compost production
- crop residues → bio-based construction materials

The rural territory becomes a network rather than a collection of isolated facilities.

Figure 3 illustrates this concept.




Figure 3. Circular rural territory and industrial symbiosis



Imagine a rural district containing a dairy farm, greenhouse complex, fruit-processing plant, wastewater treatment plant and cereal farms. Identify at least four possible circular exchanges between them. Which exchange would require the strictest quality control?

9. CIRCULAR AGRICULTURE IN CENTRAL ASIA

9.1. Regional challenges

Central Asia presents a particularly important context for circular agriculture because land, water, biomass and waste-management challenges are closely interconnected.

Important pressures include:

- water scarcity

- irrigation inefficiency
- soil salinisation and waterlogging
- wind erosion and desertification
- loss of soil organic matter
- limited rural waste collection in some areas
- uncontrolled dumping and burning
- inefficient use of crop residues
- insufficient manure management
- dependence on mineral fertilisers
- ageing rural infrastructure

The historical development of irrigation-intensive agriculture, especially for cotton and wheat, produced high levels of agricultural output but also created major environmental pressures in some areas.

The challenge is therefore not simply to introduce new waste-treatment technologies. A real circular transition must reconnect water management, land restoration, nutrient flows, biomass utilisation and rural development.

The attached reference document identifies circular opportunities in areas such as irrigation modernisation, wastewater reuse, wetland treatment, biogas, composting, legume rotations, soil-carbon restoration and rural knowledge development. It also highlights barriers including capital costs, ageing infrastructure, farmer knowledge, contamination control, technology access, financing and institutional coordination.

9.2. Circular opportunities

Several opportunities are particularly relevant.

Water efficiency and controlled reuse

Modernising irrigation infrastructure, improving scheduling and adopting drip or micro-irrigation where appropriate can increase water productivity. Treated wastewater and drainage water may provide additional resources, but salinity and contaminant risks must be managed.

Crop residue management

Straw and other crop residues can support soil protection, livestock feeding, materials or energy production. Burning residues wastes resources and creates air-pollution impacts.

The optimal solution requires determining how much biomass must remain on the land to maintain soil function and how much can be recovered sustainably.

Manure and biogas

Livestock manure offers opportunities for nutrient recycling and renewable energy. Small farm-scale digesters may be suitable in some contexts; collective facilities may offer economies of scale elsewhere.

Digestate management is crucial. A biogas system is not circular if digestate is disposed of without nutrient planning.

Soil restoration

Cover crops, crop rotations, organic amendments, agroforestry and reduced soil disturbance can help rebuild soil functions. In saline environments, these measures must be combined with appropriate irrigation and drainage management.

Rural organic waste

Source-separated organic residues from settlements and markets could potentially support composting or digestion. The main requirement is contamination control.

Mixed waste containing plastics, metals or hazardous materials cannot simply be converted into agricultural amendments.

Circular agri-food clusters

Food-processing industries can become anchor points for territorial symbiosis. Organic residues, waste heat, water and energy can be exchanged with nearby agricultural enterprises.

The principal opportunities and limitations are summarised in Table 3.

Table 3. Circular agriculture opportunities and implementation challenges in Central Asia

Domain	Circular opportunity	Main environmental benefit	Main implementation challenge
Irrigation	Efficient irrigation and controlled water reuse	Lower withdrawals and improved water productivity	Capital cost, water quality, ageing infrastructure
Nutrients	Compost, digestate, legumes and precision fertilisation	Reduced nutrient losses and external fertiliser dependence	Quality control and farmer knowledge
Crop residues	Soil return, feed, materials and bioenergy	Less burning and better biomass use	Collection and competing uses
Livestock manure	Anaerobic digestion and nutrient recovery	Renewable energy and controlled nutrient management	Finance and digestate management

Soil health	Cover crops, rotations, organic matter and agroforestry	Improved resilience and land productivity	Time horizon and incentives
Rural organics	Composting or anaerobic digestion	Reduced uncontrolled disposal	Source separation and contamination
Agri-food industries	Industrial symbiosis	Resource efficiency and reduced waste	Coordination and market development

As Table 3 shows, many of the barriers are not purely technical. Governance, investment, logistics and knowledge are equally important.

9.3. A practical transition framework

A circular transition should begin with mapping.

The first step is to identify major inputs: water, fertilisers, feed, fuel, plastics and energy.

The second is to identify outputs: food, livestock products, biomass residues, manure, wastewater, rejected products and packaging.

The third is to identify losses and environmental pressures: nitrate leaching, salinity, erosion, emissions, uncontrolled waste or unused biomass.

Only then should technologies be selected.

A practical transition pathway is shown in Figure 4.




Figure 4. Transition pathway for circular agriculture in Central Asia



Reflect

Which circular intervention would create the greatest benefit in your region: irrigation efficiency, manure management, residue recovery, composting, biogas, crop diversification or soil restoration? Explain your answer in terms of the dominant resource loss rather than the technology itself.

10. GOVERNANCE, ECONOMICS AND ENABLING CONDITIONS

Circular agriculture cannot be implemented through technology alone.

A farmer may recognise the benefits of compost but have no access to reliable quality information. A biogas facility may exist but lack a market or legal framework for digestate. Two farms may have complementary nutrient surpluses and deficits but no transport infrastructure.

Governance therefore determines whether circular opportunities become practical realities.

Important enabling conditions include:

- Clear standards. Recovered fertilisers, compost, digestate and reclaimed water need quality standards.
- Secure land tenure. Farmers are more likely to invest in soil restoration, agroforestry or long-term infrastructure when they have a sufficient planning horizon.
- Economic incentives. Subsidies, environmental taxes, payments for ecosystem services and green procurement can influence decisions.
- Spatial planning. Circular systems often benefit from locating complementary activities close together.
- Monitoring and traceability. Resource flows should be measured so that circular claims can be verified.
- Education and extension. Farmers, technicians, public authorities and businesses need practical knowledge.

Land tenure is particularly relevant. Some soil-restoration measures may take many years to show full benefits. A producer with uncertain or short-term access to land has less incentive to make long-term investments.

Circularity therefore raises a fundamental governance question:

Who invests today, who receives the future benefit and who carries the risk?



The success of a circular agricultural system may depend less on discovering a new technology than on creating agreements among existing actors who currently manage resources separately.

11. MONITORING CIRCULAR PERFORMANCE

A system should not be considered circular simply because it uses the word "circular".

Performance must be measured.

Useful indicators can be grouped into five categories.

Resource efficiency

- water use per unit of production
- mineral fertiliser input

- external feed demand
- energy use

Circular material flows

- percentage of organic residues recovered
- nutrients returned safely to land
- share of manure treated
- proportion of suitable biomass used according to cascading principles.

Soil and land condition

- soil organic carbon
- erosion
- salinity
- soil cover
- nutrient balance

Pollution prevention

- nitrate concentrations
- nutrient surplus
- greenhouse-gas emissions
- uncontrolled waste disposal
- residue burning

Socio-economic resilience

- farmer income
- local employment
- cost savings
- number of participating actors
- security of local resource supply

Table 4 proposes a simplified indicator framework.

Table 4. Indicators for monitoring circular agriculture and rural land systems

Dimension	Example indicator	Desired trend
Water	Water use per tonne of product	Decrease
Nutrients	Nitrogen surplus per hectare	Decrease
Soil	Soil organic carbon	Stable or increasing
Biomass	Share of residues recovered through safe higher-value uses	Increase
Waste	Organic waste sent to uncontrolled disposal	Decrease
Energy	Renewable energy generated from unavoidable residues	Increase where environmentally justified
Pollution	Nitrate or salinity trends in soil and groundwater	Improve or remain within safe limits
Economy	Value recovered from secondary resources	Increase
Governance	Actors participating in circular networks	Increase

As shown in Table 4, circularity must be evaluated through outcomes, not merely through activities.

A farm can recycle all manure and still pollute groundwater if application exceeds crop demand. A water-reuse system can reduce freshwater withdrawals while increasing soil salinity. A bioenergy project can recover residues but degrade soil if excessive biomass is removed.

Indicators must therefore detect trade-offs.



WARNING Maximising one circularity indicator can worsen another environmental outcome. High recycling, high water reuse or high biomass recovery should never be treated as sufficient evidence of sustainability without considering soil, water, emissions, biodiversity and resource quality.

12. CONCLUSIONS

Circular agriculture represents a transition from one-way resource consumption towards integrated management of nutrients, biomass, water, energy and rural land.

The first principle is prevention. Avoiding excessive fertiliser use, unnecessary water losses, food waste, residue burning and soil degradation is generally preferable to recovering resources after they have been lost.

The second principle is safe loop closure. Nutrients and organic matter should be returned to productive use only when their quality and environmental risks are understood.

The third principle is regeneration. A circular agricultural system must protect the land resource on which it depends. Soil organic matter, structure, biodiversity, water-holding capacity and resistance to erosion are central components of circular performance.

The fourth principle is cascading. Biomass should be kept at its highest feasible value for as long as possible. Recycling and energy recovery should not displace prevention, food use, feed use or higher-value material applications without justification.

The fifth principle is territorial integration. Farms, livestock facilities, food processors, municipalities, water managers and waste facilities may need to collaborate to close loops effectively.

For Central Asia, the transition has particular urgency because water scarcity, soil salinisation, land degradation, crop residues, manure management and rural waste are interconnected challenges. Circular agriculture offers opportunities through efficient irrigation, controlled water reuse, soil restoration, nutrient recycling, biogas, composting, crop diversification and regional resource networks.

However, circularity is not achieved by installing technologies alone. Long-term change requires quality standards, secure land management, infrastructure, financing, monitoring, farmer knowledge and cooperation among institutions.

The final lesson is therefore clear:

Circular agriculture is not the maximum recirculation of every possible material. It is the intelligent redesign of rural systems so that resources retain value, natural capital is regenerated, pollution is prevented and land remains productive and resilient over time.

13. REFERENCES

Ellen MacArthur Foundation. *Circular Economy Principles and the Biological Cycle*. Ellen MacArthur Foundation. <https://www.ellenmacarthurfoundation.org/circular-economy-principles>

European Commission (2019). *The European Green Deal*. Brussels. https://commission.europa.eu/strategy-and-policy/priorities-2019-2024/european-green-deal_en

European Commission (2020). *A Farm to Fork Strategy for a Fair, Healthy and Environmentally-Friendly Food System*. Brussels. https://knowledge4policy.ec.europa.eu/publication/communication-com2020381-farm-fork-strategy-fair-healthy-environmentally-friendly-food_en

FAO (2011). *Global Food Losses and Food Waste: Extent, Causes and Prevention*. Food and Agriculture Organization of the United Nations. <https://www.fao.org/4/mb060e/mb060e.pdf>

- FAO and ITPS (2015). *Status of the World's Soil Resources*. Food and Agriculture Organization of the United Nations and Intergovernmental Technical Panel on Soils. <https://openknowledge.fao.org/3/a-i5199e.pdf>
- FAO (2022). *FAOSTAT and Statistical Yearbook*. Food and Agriculture Organization of the United Nations. <https://openknowledge.fao.org/handle/20.500.14283/cc2211en>
- Galloway, J. N., et al. (2008). Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions. *Science*, 320, 889–892. <https://doi.org/10.1126/science.1136674>
- IPCC (2019). *Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*. <https://www.ipcc.ch/site/assets/uploads/2019/11/SRCCL-Full-Report-Compiled-191128.pdf>
- Kassam, A., Friedrich, T., and Derpsch, R. (2019). Global spread of conservation agriculture. *International Journal of Environmental Studies*. <https://doi.org/10.1080/00207233.2018.1494927>
- Ladha, J. K., et al. (2016). Global nitrogen budgets in cereals: A 50-year assessment for maize, rice and wheat production systems. *Scientific Reports*. <https://doi.org/10.1038/srep19355>
- Millennium Ecosystem Assessment (2005). *Ecosystems and Human Well-being: Synthesis*. Island Press. <https://www.millenniumassessment.org/documents/document.356.aspx.pdf>
- Newton, P., Civita, N., Frankel-Goldwater, L., Bartel, K., and Johns, C. (2020). What is regenerative agriculture? A review of scholar and practitioner definitions based on processes and outcomes. *Frontiers in Sustainable Food Systems*. <https://doi.org/10.3389/fsufs.2020.577723>
- Poeplau, C., and Don, A. (2015). Carbon sequestration in agricultural soils via cultivation of cover crops: A meta-analysis. *Agriculture, Ecosystems & Environment*, 200, 33–41. <https://doi.org/10.1016/j.agee.2014.10.024>
- United Nations Convention to Combat Desertification. *Sustainable Land Management and Land Degradation Neutrality Resources*. <https://www.unccd.int/land-and-life/land-degradation-neutrality/resources>